


PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 207/16, A61K 31/40, 31/41, C07D 207/08, 403/04	A1	(11) International Publication Number: WO 99/62880 (43) International Publication Date: 9 December 1999 (09.12.99)												
(21) International Application Number: PCT/US98/25572 (22) International Filing Date: 3 December 1998 (03.12.98) (30) Priority Data: 60/087,842 3 June 1998 (03.06.98) US (71) Applicants: GUILFORD PHARMACEUTICALS INC. [US/US]; 6611 Tributary Street, Baltimore, MD 21224 (US). AMGEN, INC. [US/US]; One Amgen Center Drive, Patent Operations M/S 27-4-A, Thousand Oaks, CA 91320-1789 (US). (72) Inventors: HAMILTON, Gregory, S.; 6501 Frederick Road, Catonsville, MD 21228 (US). NORMAN, Mark, H.; 3038 Shadow Hill Circle, Thousand Oaks, CA 91360 (US). WU, Yong-Qian; 6510 BarleyCorn Row, Columbia, MD 21044 (US). (74) Agent: NATH, Gary, M.; Nath & Associates, 6th floor, 1030 15th Street, N.W., Washington, DC 20005 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>												
(54) Title: N-LINKED SULFONAMIDES OF N-HETEROCYCLIC CARBOXYLIC ACIDS OR CARBOXYLIC ACID ISOSTERES														
(57) Abstract <p>This invention relates to novel N-linked sulfonamides of N-heterocyclic carboxylic acid and carboxylic acid isosteres, their preparation, and use for treating neurological disorders including physically damaged nerves and neurodegenerative diseases, and for treating alopecia and promoting hair growth.</p> <div data-bbox="909 1144 1299 1186">Promotion of Hair Growth by Neuroimmunophilin Ligands</div>  <table border="1"><caption>Hair Growth (Relative Index)</caption><thead><tr><th>Group</th><th>Relative Index</th></tr></thead><tbody><tr><td>Control</td><td>1.0</td></tr><tr><td>Compound 138</td><td>2.5</td></tr><tr><td>Compound 35</td><td>0.7</td></tr><tr><td>Compound 137</td><td>2.5</td></tr><tr><td>FK506</td><td>2.5</td></tr></tbody></table>			Group	Relative Index	Control	1.0	Compound 138	2.5	Compound 35	0.7	Compound 137	2.5	FK506	2.5
Group	Relative Index													
Control	1.0													
Compound 138	2.5													
Compound 35	0.7													
Compound 137	2.5													
FK506	2.5													

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

N-Linked Sulfonamides of
N-Heterocyclic Carboxylic Acids or
Carboxylic Acid Isosteres

Related Application Data

This application is a continuation-in-part of U.S. patent application serial number 60/087,842 to Hamilton et al., entitled "N-Linked Sulfonamides of N-Heterocyclic Carboxylic Acids and Carboxylic Acid Isosteres", filed June 3, 1998.

BACKGROUND OF THE INVENTION

1. Field of the Invention

This invention relates to small molecule compounds and compositions, their preparation and use for treating neurological disorders including physically damaged nerves and neurodegenerative diseases, and for treating alopecia and promoting hair growth.

2. Description of the Prior Art

It has been found that picomolar concentrations of an immunosuppressant such as FK506 and rapamycin stimulate neurite outgrowth in PC12 cells and sensory nervous, namely dorsal root ganglion cells (DRGs). Lyons et al., *Proc. of Natl. Acad. Sci.*, 1994 vol. 91, pp. 3191-3195. In whole animal experiments, FK506 has been shown to stimulate nerve regeneration following facial nerve injury and results in functional recovery in animals with sciatic nerve lesions.

Several neurotrophic factors effecting specific neuronal populations in the central nervous system have been identified. For example, it has been hypothesized that Alzheimer's disease results from a decrease or loss of nerve growth factor (NGF). It has thus been proposed

to treat Alzheimer's patients with exogenous nerve growth factor or other neurotrophic proteins such as brain derived nerve factor (BDNF), glial derived nerve factor, ciliary neurotrophic factor, and neurotrophin-3 to increase the survival of degenerating neuronal populations.

Clinical application of these proteins in various neurological disease states is hampered by difficulties in the delivery and bioavailability of large proteins to nervous system targets. By contrast, immunosuppressant drugs with neurotrophic activity are relatively small and display excellent bioavailability and specificity. However, when administered chronically, immunosuppressants exhibit a number of potentially serious side effects including nephrotoxicity, such as impairment of glomerular filtration and irreversible interstitial fibrosis (Kopp et al., 1991, J. Am. Soc. Nephrol. 1:162); neurological deficits, such as involuntary tremors, or non-specific cerebral angina such as non-localized headaches (De Groen et al., 1987, N. Engl. J. Med. 317:861); and vascular hypertension with complications resulting therefrom (Kahan et al., 1989 N. Engl. J. Med. 321: 1725).

Accordingly, there is a need for small-molecule compounds which are useful for neurotrophic effects and for treating neurodegenerative disorders.

Hair loss occurs in a variety of situations. These situations include male pattern alopecia, alopecia senilis, alopecia areata, diseases accompanied by basic skin lesions or tumors, and systematic disorders such as nutritional disorders and internal secretion disorders. The mechanisms causing hair loss are very complicated, but in some instances can be attributed to aging, genetic disposition, the activation of male hormones, the loss of blood supply to hair follicles, and scalp abnormalities.

The immunosuppressant drugs FK506, rapamycin and cyclosporin are well known as potent T-cell specific immunosuppressants, and are effective against graft rejection after organ transplantation. It has been reported that topical, but not oral, application of FK506 (Yamamoto et al., J. Invest. Dermatol., 1994, 102, 160-164; Jiang et al., J. Invest. Dermatol. 1995, 104, 523-525) and cyclosporin (Iwabuchi et al., J. Dermatol. Sci. 1995, 9, 64-69) stimulates hair growth in a dose-dependent manner. One form of hair loss, alopecia areata, is known to be associated with autoimmune activities; hence, topically administered immunomodulatory compounds are expected to demonstrate efficacy for treating that type of hair loss. The hair growth stimulating effects of FK506 have been the subject of an international patent filing covering FK506 and structures related thereto for hair growth stimulation (Honbo et al., EP 0 423 714 A2). Honbo et al. discloses the use of relatively large tricyclic compounds, known for their immunosuppressive effects, as hair revitalizing agents.

The hair growth and revitalization effects of FK506 and related agents are disclosed in many U.S. patents (Goulet et al., U.S. Patent No. 5,258,389; Luly et al., U.S. Patent No. 5,457,111; Goulet et al., U.S. Patent No. 5,532,248; Goulet et al., U.S. Patent No. 5,189,042; and Ok et al., U.S. Patent No. 5,208,241; Rupprecht et al., U.S. Patent No. 5,284,840; Organ et al., U.S. Patent No. 5,284,877). These patents claim FK506 related compounds. Although they do not claim methods of hair revitalization, they disclose the known use of FK506 for effecting hair growth. Similar to FK506 (and the claimed variations in the Honbo et al. patent), the compounds claimed in these patents are relatively large. Further, the cited patents relate to immunomodulatory compounds

for use in autoimmune related diseases, for which FK506's efficacy is well known.

Other U.S. patents disclose the use of cyclosporin and related compounds for hair revitalization (Hauer et al., U.S. Patent No. 5,342,625; Eberle, U.S. Patent No. 5,284,826; Hewitt et al., U.S. Patent No. 4,996,193). These patents also relate to compounds useful for treating autoimmune diseases and cite the known use of cyclosporin and related immunosuppressive compounds for hair growth.


However, immunosuppressive compounds by definition suppress the immune system and also exhibit other toxic side effects. Accordingly, there is a need for small molecule compounds which are useful as hair revitalizing compounds.

SUMMARY OF THE INVENTION

The present invention relates to the surprising discovery that N-heterocyclic sulfonamide compounds containing a carboxylic acid or carboxylic acid isostere moiety may be useful for treating neurodegenerative disorders and for treating alopecia and promoting hair growth. Accordingly, a novel class of sulfonamide derivatives containing an acidic moiety or an isostere thereof attached to the 2-carbon of the N-heterocyclic ring are provided. These compounds stimulate neuronal regeneration and outgrowth and as such are useful for treating neurological disorders and neurodegenerative diseases. These compounds also promote hair growth and as such are useful for treating hair loss disorders. A preferred feature of the compounds of the present invention is that they do not exert any significant immunosuppressive activity and/or are non-immunosuppressive.

A preferred embodiment of this invention is a

compound having the formula (I):



where

n is 1-3;

R₁ is selected from the group consisting of hydrogen, C₁-C₉ straight or branched chain alkyl, C₂-C₉ straight or branched chain alkenyl, aryl, heteroaryl, carbocycle, or heterocycle;

D is a bond, or a C₁-C₁₀ straight or branched chain alkyl, C₂-C₁₀ alkenyl or C₂-C₁₀ alkynyl;

R₂ is a carboxylic acid or a carboxylic acid isostere; wherein said alkyl, alkenyl, alkynyl, aryl, heteroaryl, carbocycle, heterocycle, or carboxylic acid isostere is optionally substituted with one or more substituents selected from R³, where

R³ is hydrogen, hydroxy, halo, haloalkyl, thiocarbonyl, alkoxy, alkenoxy, alkylaryloxy, aryloxy, arylalkyloxy, cyano, nitro, imino, alkylamino, aminoalkyl, sulfhydryl, thioalkyl, alkylthio, sulfonyl, C₁-C₆ straight or branched chain alkyl, C₂-C₆ straight or branched chain alkenyl or alkynyl, aryl, heteroaryl, carbocycle, heterocycle, or CO₂R⁴ where R⁴ is hydrogen or C₁-C₉ straight or branched chain alkyl or alkenyl;

or a pharmaceutically acceptable salt, ester or solvate thereof;

provided that:

when D is a bond, and R₂ is COOH,

then R₁ cannot be substituted naphthyl;

further provided that:

when D is a bond, and n is 1, and R₂ is COOH or CONHR₃,

then R₁ is not hydroxyl, methyl, ethyl, substituted or

5 unsubstituted thioethyl, benzothiazole, substituted

benzopyran, substituted benzopyrrole, substituted

benzoxazole, substituted 5-membered heterocycle

containing two N and one S heteroatoms, or substituted or

unsubstituted phenyl, phenylethyl, naphthyl, pyridyl,

10 thienyl, quinoline, tricyclic ring, aminoethyl, or

benzyl;

further provided that:

when D is a bond, and n is 2, and R₂ is COOH or

phenylbutyl ester,

15 then R₁ is not substituted phenyl, or a substituted

bicyclic ring containing two oxygen heteroatoms.

further provided that:

when D is a bond, and n is 1-2, and R₂ is a substituted or

unsubstituted carbocyclic or heterocyclic ring structure,

20 then R₁ is not substituted or unsubstituted carbocycle or

heterocycle, or hydroxy;

further provided that:

when D is a bond, and n is 1-2, and R₂ is hydroxy, alkoxy,

-SO₂(phenyl), N(R₃)₂, substituted thio or alkylthio, -NCO,

25 -PO₃(Me)₂, or -NCOOC(ethyl)phenyl,

then R₁ is not naphthalene, ethylene, substituted

tricyclic ring, or substituted or unsubstituted phenyl;

further provided that:

when D is C₁-C₃ alkyl or hexenyl, and R₂ is hydroxyl,

30 then R₁ is not substituted or unsubstituted phenyl, or

benzoimidazole;

further provided that:

when D is methyl, and n is 1, and R₂ is cyano or COOH,

then R₁ is not substituted phenyl;

35 further provided that:

when D is methyl, and n is 1, and R₂ is methoxy or N(R₃)₂,
then R₁ is not methyl, ethyl, phenylethyl, chloro
substituted alkyl, substituted oxirane, substituted
aziridine wherein one of the carbons is replaced with an
5 oxygen, substituted or unsubstituted propenyl,
substituted phenyl, benzyl, or trifluoro substituted C₂-C₃
alkyl or alkenyl;

further provided that:

when D is ethyl, and n is 2, and R₂ is hydroxyl or N(R₃)₂,
10 then R₁ is not naphthyl;

further provided that:

when D is propyl, and n is 1, and R₂ is methoxy,
then R₁ is not ethylene, cyano substituted ethyl, or
triethoxy substituted propyl;




















15 further provided that:

when D is not a bond and at least one of D and R₂ contains
at least one S or O,

then R₁ is not methyl or substituted phenyl.

A preferred embodiment of this invention is where R₂
20 is a carbocycle or heterocycle containing any combination
of CH₂, O, S, or N in any chemically stable oxidation
state, where any of the atoms of said ring structure are
optionally substituted in one or more positions with R³.

Especially preferred embodiments of this invention
25 are where R₂ is selected from the group below:



where the atoms of said ring structure may be optionally substituted at one or more positions with R³.

Another preferred embodiment of this invention is where R₂ is selected from the group consisting of -COOH, -SO₃H, -SO₂HNR³, -PO₂(R³)₂, -CN, -PO₃(R³)₂, -OR³, -SR³, -NHCOR³, -N(R³)₂, -CON(R³)₂, -CONH(O)R³, -CONHNHSO₂R³, -COHNSO₂R³, and -CONR³CN.

Preferred embodiments of this invention are compounds of the formula: (2S)-1-(phenylmethyl)sulfonyl-2-hydroxymethyl pyrrolidine; (2S)-1-(phenylmethyl)sulfonyl-2-pyrrolidinetetrazole; (2S)-1-(phenylmethyl)sulfonyl-2-pyrrolidine carbonitrile; and compounds 1-136.

Another preferred embodiment of this invention is a composition containing: a therapeutically effective amount of a compound of formula (I); a neurotrophic factor different from formula (I); and a pharmaceutically suitable carrier.

Another preferred embodiment of the invention is a method of promoting neuronal regeneration and growth in mammals, comprising administering to a mammal an effective amount of a N-linked sulfonamide of an N-heterocyclic carboxylic acid or carboxylic acid isostere.

Another preferred embodiment of the invention is a method of treating a neurological disorder in an animal, comprising administering to an animal a therapeutically effective amount of a N-linked sulfonamide of an N-heterocyclic carboxylic acid or carboxylic acid isostere to stimulate growth of damaged peripheral nerves or to promote neuronal regeneration.

Yet another preferred embodiment of the invention is a method of preventing neurodegeneration in an animal, comprising administering to an animal an effective amount of an N-linked sulfonamide of an N-heterocyclic

carboxylic acid or carboxylic acid isostere.

Yet another preferred embodiment of the invention is a method of treating alopecia or promoting hair growth in an animal, comprising administering to an animal an effective amount of an N-linked sulfonamide of an N-heterocyclic carboxylic acid or carboxylic acid isostere.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a photograph of C57 Black 6 mice before being shaved for the hair regeneration experiment.

Figure 2 is a photograph of mice treated with a vehicle after six weeks. Figure 2 shows that less than 3% of the shaved area is covered with new hair growth when the vehicle (control) is administered.

Figure 3 is a bar graph illustrating relative hair growth on shaved mice treated with N-heterocyclic carboxylic acids or carboxylic acid isosteres at 1µmole per milliliter three times per week. Hair growth was evaluated after 14 days of treatment.

DETAILED DESCRIPTION OF THE INVENTION

Definitions

"Alkyl" means a branched or unbranched saturated hydrocarbon chain comprising a designated number of carbon atoms. For example, C₁-C₆ straight or branched alkyl hydrocarbon chain contains 1 to 6 carbon atoms, and includes but is not limited to substituents such as methyl, ethyl, propyl, iso-propyl, butyl, iso-butyl, tert-butyl, n-pentyl, n-hexyl, and the like. It is also contemplated as within the scope of the present invention that "alkyl" may also refer to a hydrocarbon chain wherein any of the carbon atoms of said alkyl are optionally replaced with O, NH, S, or SO₂. For example, carbon 2 of n-pentyl can be replaced with O to form propyloxymethyl.

"Alkenyl" means a branched or unbranched unsaturated hydrocarbon chain comprising a designated number of carbon atoms. For example, C₂-C₆ straight or branched alkenyl hydrocarbon chain contains 2 to 6 carbon atoms having at least one double bond, and includes but is not limited to substituents such as ethenyl, propenyl, isopropenyl, butenyl, iso-butenyl, tert-butenyl, n-pentenyl, n-hexenyl, and the like. It is also contemplated as within the scope of the present invention that "alkenyl" may also refer to an unsaturated hydrocarbon chain wherein any of the carbon atoms of said alkenyl are optionally replaced with O, NH, S, or SO₂. For example, carbon 2 of 4-pentene can be replaced with O to form (2-propene)oxymethyl.

"Alkoxy" means the group -OR wherein R is alkyl as herein defined. Preferably, R is a branched or unbranched saturated hydrocarbon chain containing 1 to 6 carbon atoms.


The term "carbocycle" or refers to an organic cyclic moiety in which the cyclic skeleton is comprised of only carbon atoms whereas the term "heterocycle" refers to an organic cyclic moiety in which the cyclic skeleton contains one or more heteroatoms selected from nitrogen, oxygen, or sulfur and which may or may not include carbon atoms.

Thus, the term "carbocycle" refers to a carbocyclic moiety containing the indicated number of carbon atoms. The term "C₃-C₈ cycloalkyl", therefore, refers to an organic cyclic substituent in which three to eight carbon atoms form a three, four, five, six, seven, or eight-membered ring, including, for example, a cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, or cyclooctyl ring. As used herein, "carbocycle" may also refer to two or more cyclic ring systems which are fused to form, for example bicyclic, tricyclic, or other

similar bridged substituents (e.g. adamantyl).

"Aryl" refers to an aromatic carbocyclic group having a single ring, for example a phenyl ring; multiple rings, for example biphenyl; or multiple condensed rings in which at least one ring is aromatic, for example naphthyl, 1,2,3,4-tetrahydronaphthyl, anthryl, or phenanthryl, which can be unsubstituted or substituted with one or more other substituents as defined above. The substituents attached to a phenyl ring portion of an aryl moiety in the compounds of Formula (I) may be configured in the ortho-, meta-, or para- orientations.

Examples of typical aryl moieties included in the scope of the present invention may include, but are not limited to, the following:



"Aralkyl" refers to alkyl or alkylene (alkenyl) chain which is substituted with aryl, heteroaryl, carbocycle or heterocycle, or alternatively one or more aryl, heteroaryl, carbocycle, or heterocycle(s) which is/are substituted with alkyl or alkenyl, i.e.

'Alkyl/alkylene which is substituted with Ar' or 'Ar which is substituted with alkyl/alkylene'


"Heterocycle" refers to a saturated, unsaturated, or aromatic carbocyclic group having a single ring, multiple rings, or multiple condensed rings, and having at least one hetero atom such as nitrogen, oxygen, or sulfur within at least one of the rings. "Heteroaryl" refers to a heterocycle in which at least one ring is aromatic.

Any of the heterocyclic or heteroaryl groups can be unsubstituted or optionally substituted with one or more groups as defined above. Further, bi- or tri-cyclic heteroaryl moieties may comprise at least one ring which is either completely or partially saturated.

As one skilled in the art will appreciate, such heterocyclic moieties may exist in several isomeric forms, all of which are encompassed by the present invention. For example, a 1,3,5-triazine moiety is isomeric to a 1,2,4-triazine group. Such positional isomers are to be considered within the scope of the present invention. Likewise, the heterocyclic or heteroaryl groups can be bonded to other moieties in the compounds of the present invention. The point(s) of attachment to these other moieties is not to be construed as limiting on the scope of the invention. Thus, by way of example, a pyridyl moiety may be bound to other groups through the 2-, 3-, or 4-position of the pyridyl group. All such configurations are to be construed as within the scope of the present invention.

Examples of heterocyclic or heteroaryl moieties included in the scope of the present invention may

include, but are not limited to, the following:



"Halo" means at least one fluoro, chloro, bromo, or iodo moiety.

The term "pharmaceutically acceptable salt, ester, or solvate" refers to salt, ester, or solvates of the subject compounds which possess the desired pharmacological activity and which are neither biologically nor otherwise undesirable. The salt, ester, or solvates can be formed with inorganic or organic acids such as acetate, adipate, alginate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, citrate, camphorate, camphorsulfonate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, fumarate, glucoheptanoate, gluconate, glycerophosphate, hemisulfate, heptanoate, hexanoate, hydrochloride hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, lactate, maleate, methanesulfonate, naphthylate, 2-naphthalenesulfonate, nicotinate, oxalate, sulfate, thiocyanate, tosylate and undecanoate. Base salt, ester, or solvates include ammonium salts, alkali metal salts such as lithium, sodium and potassium salts, alkaline earth metal salts such as calcium and magnesium salts, salt with organic bases such as dicyclohexylamine salts, N-methyl-D-glucamine, and salts with amino acids such as arginine, lysine, and so forth. Also, the basic nitrogen-containing groups can be quarternized with such agents as: 1) lower alkyl halides, such as methyl, ethyl, propyl, and butyl chloride, bromides and iodides; 2) dialkyl sulfates like dimethyl, diethyl, dibutyl and diamyl sulfates; 3) long chain alkyls such as decyl, lauryl, myristyl and stearyl substituted with one or more halide such as chloride, bromide and iodide; and 4) aryl or arylalkyl halides like benzyl and phenethyl bromide and others.

The compounds of this invention may possess at least one asymmetric center and thus can be produced as

mixtures of stereoisomers or as individual enantiomers or diastereomers. The individual stereoisomers may be obtained by using an optically active starting material, by resolving a racemic or non-racemic mixture of an intermediate at some appropriate stage of the synthesis, or by resolution of the compound of formula (I). It is understood that the individual stereoisomers as well as mixtures (racemic and non-racemic) of stereoisomers are encompassed by the scope of the present invention. The S-stereoisomer at atom 1 of formula I is a most preferred embodiment of the invention.

"Stereoisomers" are isomers that differ only in the way the atoms are arranged in space.

"Isomers" are different compounds that have the same molecular formula and includes cyclic isomers such as (iso)indole and other isomeric forms of cyclic moieties.

"Enantiomers" are a pair of stereoisomers that are non-superimposable mirror images of each other.

"Diastereoisomers" are stereoisomers which are not mirror images of each other.


"Racemic mixture" means a mixture containing equal parts of individual enantiomers. "Non-racemic mixture" is a mixture containing unequal parts of individual enantiomers or stereoisomers.

"Isosteres" are different compounds that have different molecular formulae but exhibit the same or similar properties. For example, tetrazole is an isostere of carboxylic acid because it mimics the properties of carboxylic acid even though they both have very different molecular formulae. Tetrazole is one of many possible isosteric replacements for carboxylic acid. Other carboxylic acid isosteres contemplated by the present invention include

-COOH, -SO₃H, -SO₂HNR³, -PO₂(R³)₂, -CN, -PO₃(R³)₂, -OR³, -SR³, -NHCOR³, -N(R³)₂, -CON(R³)₂, -CONH(O)R³, -CONHNHSO₂R³,

-COHNSO₂R³, and -CONR³CN.

In addition, carboxylic acid isosteres can include 5-7 membered carbocycles or heterocycles containing any combination of CH₂, O, S, or N in any chemically stable oxidation state, where any of the atoms of said ring structure are optionally substituted in one or more positions. The following structures are non-limiting examples of preferred carbocyclic and heterocyclic isosteres contemplated by this invention.



where the atoms of said ring structure may be optionally substituted at one or more positions with R^3 . The present invention contemplates that when chemical substituents are added to a carboxylic isostere then the inventive compound retains the properties of a carboxylic isostere. The present invention contemplates that when a carboxylic isostere is optionally substituted with one or more moieties selected from R^3 , then the substitution can not eliminate the carboxylic acid isosteric properties of the inventive compound. The present invention contemplates that the placement of one or more R^3 substituents upon a carbocyclic or heterocyclic carboxylic acid isostere shall not be at an atom(s) which maintains or is integral to the carboxylic acid isosteric properties of the inventive compound if such a substituent(s) would destroy the carboxylic acid isosteric properties of the inventive compound.

Other carboxylic acid isosteres not specifically exemplified or described in this specification are also contemplated by the present invention.

The term "preventing neurodegeneration" as used herein includes the ability to inhibit or prevent neurodegeneration in patients newly diagnosed as having a neurodegenerative disease, or at risk of developing a new degenerative disease and for inhibiting or preventing further neurodegeneration in patients who are already suffering from or have symptoms of a neurodegenerative disease when the compounds are given concurrently.

The term "treatment" as used herein covers any treatment of a disease and/or condition in an animal, particularly a human, and includes:

(i) preventing a disease and/or condition from occurring in a subject which may be predisposed to the disease and/or condition but has not yet been diagnosed as having it;

(ii) inhibiting the disease and/or condition, i.e., arresting its development; or

(iii) relieving the disease and/or condition, i.e., causing regression of the disease and/or condition.

5 The system used in naming the compounds of the present invention is shown below, using a compound of formula I as an example.

A compound of the present invention, especially formula I, wherein n is 1, D is a bond, R₁ is
10 phenylmethyl, and R₂ is -CN, is named (2S)-1-(phenylmethyl) sulfonyl-2-pyrrolidine carbonitrile.

"Alopecia" refers to deficient hair growth and partial or complete loss of hair, including without limitation androgenic alopecia (male pattern baldness),
15 toxic alopecia, alopecia senilis, alopecia areata, alopecia pelada and trichotillomania. Alopecia results when the pilar cycle is disturbed. The most frequent phenomenon is a shortening of the hair growth or anagen phase due to cessation of cell proliferation. This
20 results in an early onset of the catagen phase, and consequently a large number of hairs in the telogen phase during which the follicles are detached from the dermal papillae, and the hairs fall out. Alopecia has a number of etiologies, including genetic factors, aging, local
25 and systemic diseases, febrile conditions, mental stresses, hormonal problems, and secondary effects of drugs.

"Pilar cycle" refers to the life cycle of hair follicles, and includes three phases:

- 30 (1) the anagen phase, the period of active hair growth which, insofar as scalp hair is concerned, lasts about three to five years;
- (2) the catagen phase, the period when growth stops and the follicle atrophies which, insofar as
35 scalp hair is concerned, lasts about one to two

weeks; and

- (3) the telogen phase, the rest period when hair progressively separates and finally falls out which, insofar as scalp hair is concerned, lasts about three to four months.

Normally 80 to 90 percent of the follicles are in the anagen phase, less than 1 percent being in the catagen phase, and the rest being in the telogen phase. In the telogen phase, hair is uniform in diameter with a slightly bulbous, non-pigmented root. By contrast, in the anagen phase, hair has a large colored bulb at its root.

"Promoting hair growth" refers to maintaining, inducing, stimulating, accelerating, or revitalizing the germination of hair.

"Treating alopecia" refers to:

- (i) preventing alopecia in an animal which may be predisposed to alopecia; and/or
- (ii) inhibiting, retarding or reducing alopecia; and/or
- (iii) promoting hair growth; and/or
- (iv) prolonging the anagen phase of the hair cycle; and/or

(v) converting vellus hair to growth as terminal hair. Terminal hair is coarse, pigmented, long hair in which the bulb of the hair follicle is seated deep in the dermis. Vellus hair, on the other hand, is fine, thin, non-pigmented short hair in which the hair bulb is located superficially in the dermis. As alopecia progresses, the hairs change from the terminal to the vellus type.

The term "neurotrophic" as used herein includes without limitation the ability to stimulate neuronal regeneration or growth and/or the ability to prevent or treat neurodegeneration.

The term "non-immunosuppressive" refers to the inability of the compounds of the present invention to trigger an immune response when compared to a control such as FK506 or cyclosporin A. Assays for determining immunosuppression are well known to those of ordinary skill in the art. Specific non-limiting examples of well known assays include PMA and OKT3 assays wherein mitogens are used to stimulate proliferation of human peripheral blood lymphocytes (PBC). Compounds added to such assay systems are evaluated for their ability to inhibit such proliferation.

Compounds of the Invention

The present invention relates to the surprising discovery that N-linked sulfonamides of N-heterocyclic carboxylic acid or carboxylic acid isostere compounds are neurotrophic and are able to treat alopecia. Accordingly, a novel class of compounds are provided. A preferred feature of the compounds of the present invention is that they do not exert any significant immunosuppressive activity.

Preferred compounds of the present invention contain carboxylic acid moieties and other isosteric replacements for carboxylic acid moieties, of which several examples are specified herein. Other isosteric replacements for carboxylic acid moieties, known to those skilled in the art of medicinal chemistry, are within the scope of the invention if not otherwise specified.

The neurotrophic compounds of this invention can be periodically administered to a patient undergoing treatment for neurological disorders or for other reasons in which it is desirable to stimulate neuronal regeneration and growth, such as in various peripheral neuropathic and neurological disorders relating to neurodegeneration. The compounds of this invention can also be administered to mammals other than humans for

treatment of various mammalian neurological disorders.

The novel compounds of the present invention possess an excellent degree of neurotrophic activity. This activity is useful in the stimulation of damaged neurons, the promotion of neuronal regeneration, the prevention of neurodegeneration, and in the treatment of several neurological disorders known to be associated with neuronal degeneration and peripheral neuropathies. The neurological disorders that may be treated include but are not limited to: trigeminal neuralgia, glossopharyngeal neuralgia, Bell's Palsy, myasthenia gravis, muscular dystrophy, amyotrophic lateral sclerosis, progressive muscular atrophy, progressive bulbar inherited muscular atrophy, herniated, ruptured or prolapsed invertebrate disk syndromes, cervical spondylosis, plexus disorders, thoracic outlet destruction syndromes, peripheral neuropathic such as those caused by lead, dapsone, ticks, porphyria, or Gullain-Barré syndrome, Alzheimer's disease, and Parkinson's disease.

The above discussion relating to the utility and administration of the compounds of the present invention also applies to the pharmaceutical compositions of the present invention.

The term "pharmaceutically acceptable carrier" as used herein refers to any carrier, diluent, excipient, suspending agent, lubricating agent, adjuvant, vehicle, delivery system, emulsifier, disintegrant, absorbant, preservative, surfactant, colorant, flavorant, or sweetener.

For these purposes the compounds of the present invention may be administered orally, parenterally, by inhalation spray, topically, rectally, nasally, buccally, vaginally or via an implanted reservoir in dosage formulations containing conventional non-toxic

pharmaceutically-acceptable carriers, adjuvants and vehicles. The term parenteral as used herein includes subcutaneous, intravenous, intramuscular, intraperitoneally, intrathecally, intraventricularly, intrasternal and intracranial injection or infusion techniques.

For oral administration, the compounds of the present invention may be provided in any suitable dosage form known in the art. For example, the compositions may be incorporated into tablets, powders, granules, beads, chewable lozenges, capsules, liquids, aqueous suspensions or solutions, or similar dosage forms, using conventional equipment and techniques known in the art. Tablet dosage forms are preferred. Tablets may contain carriers such as lactose and corn starch, and/or lubricating agents such as magnesium stearate. Capsules may contain diluents including lactose and dried corn starch. Aqueous suspensions may contain emulsifying and suspending agents combined with the active ingredient.

When preparing dosage form incorporating the compositions of the invention, the compounds may also be blended with conventional excipients such as binders, including gelatin, pregelatinized starch, and the like; lubricants, such as hydrogenated vegetable oil, stearic acid, and the like; diluents, such as lactose, mannose, and sucrose; disintegrants, such as carboxymethylcellulose and sodium starch glycolate; suspending agents, such as povidone, polyvinyl alcohol, and the like; absorbants, such as silicon dioxide; preservatives, such as methylparaben, propylparaben, and sodium benzoate; surfactants, such as sodium lauryl sulfate, polysorbate 80, and the like; colorants such as F.D. & C. dyes and lakes; flavorants; and sweeteners.

Compositions and methods of the invention also may utilize controlled release technology. Thus, for

example, the inventive compounds may be incorporated into a hydrophobic polymer matrix for controlled release over a period of days. Such controlled release films are well known to the art. Particularly preferred are transdermal delivery systems. Other examples of polymers commonly employed for this purpose that may be used in the present invention include nondegradable ethylene-vinyl acetate copolymer and degradable lactic acid-glycolic acid copolymers which may be used externally or internally. Certain hydrogels such as poly(hydroxyethylmethacrylate) or poly(vinylalcohol) also may be useful, but for shorter release cycles than the other polymer releases systems, such as those mentioned above.

To be effective therapeutically as central nervous system targets, the compounds of the present invention should readily penetrate the blood-brain barrier when peripherally administered. Compounds which cannot penetrate the blood-brain barrier can be effectively administered by an intraventricular route or other appropriate delivery system suitable for administration to the brain.

The compounds of the present invention may be administered in the form of sterile injectable preparations, for example, as sterile injectable aqueous or oleaginous suspensions. These suspensions may be formulated according to techniques known in the art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparations may also be sterile injectable solutions or suspensions in non-toxic parenterally-acceptable diluents or solvents, for example, as solutions in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as solvents or suspending

mediums. For this purpose, any bland fixed oil may be employed including synthetic mono- or di-glycerides. Fatty acids such as oleic acid and its glyceride derivatives, including olive oil and castor oil, especially in their polyoxyethylated versions, are useful in the preparation of injectables. These oil solutions or suspensions may also contain long-chain alcohol diluents or dispersants.

The compounds of this invention may also be administered rectally in the form of suppositories. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient which is solid at room temperature, but liquid at rectal temperature and, therefore, will melt in the rectum to release the drug. Such materials include cocoa butter, beeswax and polyethylene glycols.

The compounds of this invention may also be administered topically, especially when the conditions addressed for treatment involve areas or organs readily accessible by topical application, including neurological disorders of the eye, the skin, or the lower intestinal tract. Suitable topical formulations are readily prepared for each of these areas.

For topical application to the eye, or ophthalmic use, the compounds can be formulated as micronized suspensions in isotonic, pH adjusted sterile saline, or, preferably, as solutions in isotonic, pH adjusted sterile saline, either with or without a preservative such as benzylalkonium chloride. Alternatively for the ophthalmic uses the compounds may be formulated in an ointment such as petrolatum.

For topical application to the skin, the compounds can be formulated in a suitable ointment containing the compound suspended or dissolved in, for example, a mixture with one or more of the following: mineral oil,

liquid petrolatum, white petrolatum, propylene glycol, polyoxyethylene polyoxypropylene compound, emulsifying wax and water. Alternatively, the compounds can be formulated in a suitable lotion or cream containing the active compound suspended or dissolved in, for example, a mixture of one or more of the following: mineral oil, sorbitan monostearate, polysorbate 60, cetyl esters wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water.

Topical application for the lower intestinal tract can be effected in a rectal suppository formulation (see above) or in a suitable enema formulation.

Dosage levels on the order of about 0.1 mg to about 10,000 mg of the active ingredient compound are useful in the treatment of the above conditions, with preferred levels of about 0.1 mg to about 1,000 mg. The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. Typically, *in vitro* dosage-effect results provide useful guidance on the proper doses for patient administration. Studies in animal models are also helpful. The considerations for determining the proper dose levels are well known in the art.

It is understood, however, that a specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, rate of excretion, drug combination, and the severity of the particular disease being treated and form of administration.

To effectively treat alopecia or promote hair growth, the compounds used in the inventive methods and pharmaceutical compositions must readily affect the targeted areas. For these purposes, the compounds are


preferably administered topically to the skin.

For topical application to the skin, the compounds can be formulated into suitable ointments containing the compounds suspended or dissolved in, for example, mixtures with one or more of the following: mineral oil, liquid petrolatum, white petrolatum, propylene glycol, polyoxyethylene polyoxypropylene compound, emulsifying wax and water. Alternatively, the compounds can be formulated into suitable lotions or creams containing the active compound suspended or dissolved in, for example, a mixture of one or more of the following: mineral oil, sorbitan monostearate, polysorbate 60, cetyl ester wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water.

The compounds can be administered for treatment of hair loss with other hair revitalizing agents. Specific dose levels for the other hair revitalizing agents will depend upon the factors previously stated and the effectiveness of the drug combination.




Other routes of administration known in the pharmaceutical art are also contemplated by this invention.

Specific embodiments of the inventive compounds are presented in Table I. The present invention contemplates employing the compounds of Table I, below, for use in compositions and methods to prevent and/or treat a neurological disorder in an animal, and for use in compositions and methods to treat alopecia and promote hair growth in an animal, and all other uses suggested in this specification.


TABLE I

	No.	n	D	R2	R1
5	1	1	bond	COOH	Benzyl
	2	1	bond	COOH	α -MethylBenzyl
	3	1	bond	COOH	4-MethylBenzyl
	4	1	bond	Tetrazole	Benzyl
	5	1	bond	SO ₃ H	α -MethylBenzyl
10	6	1	CH ₂	COOH	4-MethylBenzyl
	7	1	bond	SO ₂ HNMe	Benzyl
	8	1	bond	CN	α -MethylBenzyl
	No.	n	D	R2	R1
15	9	1	bond	PO ₃ H ₂	4-MethylBenzyl
	10	2	bond	COOH	Benzyl
	11	2	bond	COOH	α -MethylBenzyl
	12	2	bond	COOH	4-MethylBenzyl
	13	2	bond	COOH	3,4,5-trimethoxy phenyl
20	14	2	bond	COOH	Cyclohexyl
	15	2	bond	PO ₂ HEt	i-propyl
	16	2	bond	PO ₃ HPropyl	ethyl
	17	2	bond	PO ₃ (Et) ₂	Methyl
	18	2	bond	OMe	tert-butyl
25	19	2	bond	OEt	n-pentyl
	20	2	bond	OPropyl	n-hexyl
	21	1	bond	OButyl	Cyclohexyl
	22	1	bond	OPentyl	cyclopentyl
	23	1	bond	OHexyl	n-heptyl

	24	1	bond	SMe	n-octyl
	25	1	bond	SEt	n-nonyl
	26	2	bond	SPropyl	2-indolyl
	27	2	bond	SButyl	2-furyl
5	28	2	bond	NHCOMe	2-thiazolyl
	29	2	bond	NHCOEt	2-thienyl
	30	1	CH ₂	N(Me) ₂	2-pyridyl
	31	1	(CH ₂) ₂	N(Me)Et	benzyl
	32	1	(CH ₂) ₃	CON(Me) ₂	benzyl
10	33	1	(CH ₂) ₄	CONHMe	benzyl
	34	1	(CH ₂) ₅	CONHEt	benzyl
	35	1	(CH ₂) ₆	CONHPropyl	1,1-dimethylpropyl
	36	1	bond	CONH(O)Me	Benzyl
	37	1	bond	CONH(O)Et	α-Methylphenyl
15	38	1	bond	CONH(O)Propyl	4-Methylphenyl
	39	2	bond	COOH	Benzyl
	40	2	bond	COOH	α-Methylphenyl
	41	2	bond	COOH	4-Methylphenyl
	No.	n	D	R2	R1
20	42	1	CH ₂	COOH	benzyl
	43	1	(CH ₂) ₂	COOH	benzyl
	44	1	(CH ₂) ₃	COOH	benzyl
	45	1	(CH ₂) ₄	COOH	benzyl
	46	1	(CH ₂) ₅	COOH	benzyl
25	47	1	(CH ₂) ₆	COOH	benzyl
	48	1	(CH ₂) ₇	COOH	benzyl
	49	1	(CH ₂) ₈	COOH	benzyl
	50	1	(CH ₂) ₉	COOH	benzyl
	51	1	(CH ₂) ₁₀	COOH	benzyl
30	52	1	C ₂ H ₂	COOH	benzyl
	53	1	2-OH, Et	COOH	benzyl
	54	1	2butylene	COOH	benzyl
	55	1	i-Pro	COOH	benzyl
	56	1	tert-Bu	COOH	benzyl
35	57	1	2-nitro	COOH	benzyl


			Hexyl		
	58	3	(CH ₂) ₂	CN	benzyl
	59	1	(CH ₂) ₃	CN	benzyl
	60	3	bond	CONHNHSO ₂ Me	Benzyl
5	61	3	bond	CONHNHSO ₂ Et	α-Methylphenyl
	62	3	bond	CONHSO ₂ Me	4-Methylphenyl
	63	2	bond	CONHNHSO ₂ Et	Phenyl
	64	2	bond	CON(Me)CN	α-Methylphenyl
	65	2	bond	CON(Et)CN	4-Methylphenyl
10	66	1	(CH ₂) ₂	COOH	methyl
	67	1	(CH ₂) ₃	COOH	ethyl
	68	1	(CH ₂) ₄	COOH	n-propyl
	69	1	(CH ₂) ₅	COOH	t-butyl
	70	1	(CH ₂) ₆	COOH	Pentyl
15	71	1	(CH ₂) ₇	COOH	Hexyl
	72	1	(CH ₂) ₈	COOH	Septyl
	73	1	(CH ₂) ₉	COOH	Octyl
	74	1	(CH ₂) ₁₀	COOH	Nonyl
	No.	n	D	R2	R1
20	75	1	C ₂ H ₂	COOH	Cyclohexyl
	76	1	bond		benzyl
	77	1	bond		benzyl
	78	1	bond		benzyl

79 1 bond




benzyl

80 1 bond



benzyl

81 1 bond




benzyl

No. n D
82 1 bond


R2

R1

benzyl




83 1 bond




benzyl

84 1 bond




benzyl

85 1 bond



benzyl

86 1 bond



benzyl


5

No. n D
87 1 bond


R2

R1

benzyl




88 1 bond




benzyl

89 1 bond




benzyl

90 1 bond



benzyl

91 1 bond



benzyl


5

No. n D
92 1 bond

R2


R1

benzyl



10


93 1 bond



benzyl

94 1 bond

benzyl




95	1	bond	CH ₂ OH	benzyl
96	1	bond	CONH ₂	benzyl
97	1	bond	CN	benzyl

5

Additional claimed or comparative carboxylic acids and isosteres of N-heterocyclic compounds which also show the remarkable neurotrophic and hair growth effects of the present invention are shown below in

10 Table II:

TABLE II



	Cpd	n	D	R ₂	L	R ₁
	A	1	bond	COOH	SO ₂	Benzyl
5	B	1	bond	CONH ₂	SO ₂	Benzyl
	C	1	bond	-CN	SO ₂	Benzyl
	D	1	bond	tetrazole	SO ₂	Benzyl
	E	1	CH ₂	-OH	SO ₂	Benzyl
	F	1	bond	COOH	1,2-dioxoethyl	1,1-dimethylpropyl
10	G	2	bond	COOH	1,2-dioxoethyl	1,1-dimethylpropyl
	H	1	CH ₂	OH	1,2-dioxoethyl	1,1-dimethylpropyl
	I	1	bond	tetrazole	1,2-dioxoethyl	1,1-dimethylpropyl
	J	1	bond	-CN	1,2-dioxoethyl	1,1-dimethylpropyl
	K	2	bond	CONH ₂	1,2-dioxoethyl	1,1-dimethylpropyl
15	where Y and Z are both carbon for compounds A-K,					
	L	1	bond	COOH	1,2-dioxoethyl	1,1-dimethylpropyl
	M	1	bond	COOH	1,2-dioxoethyl	1,1-dimethylpropyl
	where Z is S for compound H or					
20	where Y is S for compound I.					

Pharmaceutical Compositions of the Present Invention

The present invention relates to a pharmaceutical composition comprising:

- 25 (i) an effective amount of an N-linked sulfonamide of N-heterocyclic carboxylic acid or carboxylic acid isostere compound; and
- (ii) a pharmaceutically acceptable carrier.

The present invention also relates to a pharmaceutical composition comprising:

- (i) an effective amount of an N-linked sulfonamide of N-heterocyclic carboxylic acid or carboxylic acid isostere compound for treating neurodegenerative diseases, neurological disorders, and nerve damage, or promoting nerve growth in an animal; and
- (ii) a pharmaceutically acceptable carrier.

The present invention also relates to a pharmaceutical composition comprising:

- (i) an effective amount of an N-linked sulfonamide of N-heterocyclic carboxylic acid or carboxylic acid isostere compound for treating alopecia or promoting hair growth in an animal; and
- (ii) a pharmaceutically acceptable carrier.

Neurotrophic compounds can be administered with other neurotrophic agents such as neurotrophic growth factor, brain derived growth factor, glial derived growth factor, ciliary neurotrophic factor, insulin growth factor and active truncated derivatives thereof, acidic fibroblast growth factor, basic fibroblast growth factor, platelet-derived growth factors, neurotrophin-3 and neurotrophin 4/5. The dosage level of other neurotrophic drugs will depend upon the factors previously stated and the neurotrophic effectiveness of the drug combination.

Methods of the Present Invention

The present invention relates to the use of any of the compounds seen in Table I and II and other compounds embodied herein, in the preparation of a medicament for the treatment of a disease such as peripheral neuropathy caused by physical injury or disease state, physical damage to the brain, physical

damage to the spinal cord, stroke associated with brain damage, Alzheimer's Disease, Parkinson's Disease, and amyotrophic lateral sclerosis. The present invention also relates to the use of

5 carboxylic acid and carboxylic acid isostere compounds for treating the above-mentioned neuropathies, neurological disorders, and neurological damage.

The present invention also relates to a method for treating alopecia or promoting hair growth in an animal, which comprises administering to said animal

10 an effective amount of, an N-linked sulfonamide of an N-heterocyclic carboxylic acid or carboxylic acid isostere. The present invention also relates to using the inventive compounds and compositions in the

15 preparation of a medicament for the treatment of alopecia or promoting hair growth in an animal.

The inventive method is particularly useful for treating male pattern alopecia, alopecia senilis, alopecia areata, alopecia resulting from skin lesions

20 or tumors, alopecia resulting from cancer therapy such as chemotherapy and radiation, and alopecia resulting from systematic disorders such as nutritional disorders and internal secretion disorders.

It is understood, however, that a specific dose

25 level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, rate of excretion, drug combination, and the severity

30 of the particular disease or disorder being treated and form of administration.

MPTP Model of Parkinson's Disease in Mice

MPTP lesioning of dopaminergic neurons in mice is

35 used as an animal model of Parkinson's Disease. Four

week old male CD1 white mice are dosed i.p. with 30 mg/kg of MPTP for 5 days. Test compounds (4 mg/kg), or vehicle, are administered s.c. along with the MPTP for 5 days, as well as for an additional 5 days following cessation of MPTP treatment. At 18 days following MPTP treatment, the animals are sacrificed and the striata dissected and perfusion-fixed. Immunostaining is performed on sagittal and coronal brain sections using anti-tyrosine hydroxylase 1 g to quantitate survival and recovery of dopaminergic neurons. In animals treated with MPTP and vehicle, a substantial loss of functional dopaminergic terminals is observed as compared to non-lesioned animals. Lesioned animals receiving test compounds show a significant recovery of TH-stained dopaminergic neurons. This model presents quantitation for the recovery of TH-positive dopaminergic neurons in the striatum of animals receiving the compounds of the present invention. Table III presents the percent recovery of dopaminergic neurons in the first (concurrent dosing) paradigm in animals receiving Compound 1, (2S)-1-[(phenylmethyl)sulfonyl]-2-pyrrolidinecarboxylic acid as well related compounds of the present invention.

Table III, below, shows the remarkable neuroregenerative effects of carboxylic acid or carboxylic acid isostere related compounds illustrating the neurotrophic capability of carboxylic acid isosteres as a class showing that lesioned animals receiving the carboxylic acid or carboxylic acid isostere compounds provide a remarkable recovery of TH-stained dopaminergic neurons.

Table III - MPTP Neurodegenerative Model

		% Recovery
	Compound A	24.4 %
	Cmpds B-E	ND
5	Compound F	26.7 %
	Compound G	ND
	Compound H	23.2 %
	Compound I	19.6 %
	Compound J	34.1 %
10	Compound K	46.5 %
	Compound L	14.0 %
	Compound M	ND

Percent striatal innervation density was quantitated in brain sections with an anti-tyrosine hydroxylase immunoglobulin, which is indicative of functional dopaminergic neurons. The striatal innervation density of 23% for animals pretreated with only a vehicle and administered a vehicle orally during treatment, is indicative of normal non-lesioned striatal tissue. Striatal innervation density is reduced to 5% for animals pretreated with MPTP and administered a vehicle orally during treatment, and is indicative of MPTP-induced lesioning. Surprisingly, striatal innervation density is increased 8-13% for animals pretreated with MPTP and administered 0.4 mg/kg of compound orally during treatment, indicating substantial neuronal regeneration after induction of MPTP-derived lesions.

In Vivo Hair Generation Test With C57 Black 6 Mice

C57 black 6 mice are used to demonstrate the hair revitalizing properties of the ureas and carbamates of N-heterocyclic carboxylic acids or carboxylic acid isosteres. Referring now to FIGS. 1 and 2 of the

drawings, C57 black 6 mice, approximately 7 weeks old, had an area of about 2 inches by 2 inches on their hindquarters shaved to remove all existing hair. Care was taken not to nick or cause abrasion to the underlaying dermal layers. The animals were in anagen growth phase, as indicated by the pinkish color of the skin. Referring now to FIG. 2, four animals per group were treated by topical administration with 20% propylene glycol vehicle (FIG. 2), or related compounds dissolved in the vehicle. The animals were treated with vehicle or N-heterocyclic carboxylic acids or isosteres every 48 hours (3 applications total over the course of 5 days) and the hair growth was allowed to proceed for 6 weeks. Hair growth was quantitated by the percent of shaved area covered by new hair growth during this time period.

FIG. 2 shows that animals treated with vehicle exhibited only a small amount of hair growth in patches or tufts, with less than 3% of the shaved area covered with new growth.

In contrast, FIG. 3 shows that animals treated for 2 weeks with the N-heterocyclic carboxylic acid compounds i.e. compound F, compound G, and compound K exhibited dramatic hair growth, covering greater than 25% of the shaved area in all animals for two of the compounds.

FIG. 3 shows the relative hair growth on shaven C57 black 6 mice 14 days after being treated with N-heterocyclic carboxylic acids or carboxylic acid isosteres. The mice had a 2 x 2 inch region on their backside shaved to remove all hair. Care was taken not to nick or cause abrasion to the underlying dermal layers. Compounds at a concentration of 1 μ mole per milliliter were carefully applied to the shaved area of the mice (5 mice per group) three times per week.

Hair growth was evaluated 14 days after initiation of drug treatment. The relative scale for assessing hair growth is as follows:

0 = no growth;

5 1 = beginning of growth in small tufts;

2 = hair growth covering over <25% of shaved area;

3 = hair growth covering over >25% of shaved area, but less than 50% of shaved area;

10 4 = hair growth covering over >50% of shaved area, but less than 75% of shaved area;


5 = complete hair growth of shaved area.

15 The following examples are illustrative of preferred embodiments of the invention and are not to be construed as limiting the invention thereto. All polymer molecular weights are mean average molecular weights. All percentages are based on the percent by weight of the final delivery system or formulation prepared unless otherwise indicated and all totals equal 100% by weight.


20


EXAMPLES

25 The inventive compounds may be prepared by a variety of synthetic sequences that utilize established chemical transformations. An exemplary general pathway to the present compounds is described in Scheme I, Scheme II, and Scheme III.

SCHEME I

5

SCHEME II

SCHEME IIIEXAMPLE 1

Synthesis of (2S)-N-(benzylsulfonyl)-2-pyrrolidinecarboxylic acid (Table I Compound 1) (Table II and Scheme I Compound A)

To a cooled (0°C) solution of proline methyl ester hydrochloride salt (5.0 g; 30.19 mmol) in 200 mL of methylene chloride was added triethylamine (35mL) and benzenesulfonyl chloride (5.75 g; 30.19 mmol). The mixture was stirred for one hour at 0°C and then washed with 2 x 100 mL of water. The organic phase was dried and concentrated. Chromatography eluting with 50% EtOAc/hexane delivered 8.14 g (5%) of the N-sulfonamide methyl ester, which was dissolved in 120 mL of methanol, cooled to 0°C , and treated with 40 mL of 1 N lithium hydroxide. The mixture was stirred for 1 hour at 0°C and then overnight at room temperature. After making the reaction mixture acidic (pH 1) with 1 N HCl, the product was extracted into methylene chloride and dried and concentrated to yield 4.25 g of (2S)-N-(benzylsulfonyl)-2-pyrrolidinecarboxylic acid (A) as a white solid, ^1H NMR (CDCl_3 , 400 MHz): δ 1.85-1.90 (m, 2H); 2.08 (m, 1H); 2.18 (m, 1H); 3.04 (m, 1H); 3.27 (m, 1H); 4.32-4.35 (m, 2H); 4.45 (m, 1H); 4.45 (m, 2H); 7.36 (m, 3H); 7.48 (m, 2H); 10.98 (br, 1H).

EXAMPLE 2

Synthesis of (2S)-1-(phenylmethylsulfonyl)-2-hydroxymethyl pyrrolidine (Compound 95) (Scheme III Compound E).

5 To a solution of (S)-(+)-2-pyrrolidinemethanol (1.01 g, 10 mmol) and triethylamine (1.5 ml, 11 mmol) in 30 ml methylene chloride was added 1.9 g (10 mmol) α -toluenesulfonyl chloride at 0°C with stirring. The reaction was gradually warmed up to room temperature and stirred overnight. The mixture was diluted with water, and extracted into 200 ml methylene chloride. The organic extract was concentrated and further purified by silica gel to give 1.5 g product as a white solid (58.9% yield). ¹H NMR (CDCl₃): d 0.71-1.88 (m, 4H); 2.05 (br, 1H, OH); 3.22 (m, 2H); 3.47 (m, 2H); 3.67 (m, 1H); 4.35 (s, 2H); 7.26-7.44 (m, 5H, aromatic).

EXAMPLE 3

20 Synthesis of (2S)-1-(phenylmethyl)sulfonyl-2-pyrrolidinecarboxamide (Compound 96) (Scheme II Compound B).

To a solution of L-prolinamide (2.28 g, 20 mmol) and triethylamine (5.76 ml, 42 mmol) in 40 ml methylene chloride was added 3.92 g (20 mmol) α -toluenesulfonyl chloride at 0°C with stirring. The reaction was gradually warmed up to room temperature and stirred overnight. The mixture was diluted with water, and extracted into 200 ml methylene chloride. The organic extract was concentrated and further purified by silica gel to give 3.0 g product as a white solid (55.7% yield). ¹H NMR (CDCl₃): d 0.89 (m, 3H); 2.25 (m, 1H); 3.40 (m, 1H); 3.50 (m, 1H); 3.96 (m, 1H); 4.35 (s, 2H); 7.39-7.45 (m, 5H, aromatic).

EXAMPLE 4

Synthesis of (2S)-1-(phenylmethyl)sulfonyl-2-pyrrolidinecarbonitrile (Compound 97) (Scheme II Compound C).

To a solution of 0.67 ml DMF (8.7 mmol) in 10 ml acetonitrile at 0°C was added 0.70 ml (8.0 mmol) oxalyl chloride. A white precipitate was formed immediately and was accompanied by gas evolution. When complete, a solution of 2.0 g (7.5 mmol) of (2S)-1-(phenylmethyl)sulfonyl-2-pyrrolidinecarboxamide in 5.0 ml acetonitrile was added. When the mixture became homogeneous, 1.35 ml (16.5 mmol) pyridine was added. After 5 min., the mixture was diluted with water, and extracted by 200 ml ethyl acetate. The organic layer was concentrated and further purified by silica gel to give 1.5 g product as a white solid (80% yield). ¹H NMR (CDCl₃): δ 01.92 (m, 2H); 2.01 (m, 1H); 2.11 (m, 1H); 3.45 (m, 2H); 4.35 (s, 2H); 4.65 (m, 1H); 7.26-7.45 (m, 5H, aromatic).

EXAMPLE 5

Synthesis of (2S)-1-(phenylmethyl)sulfonyl-2-pyrrolidinetetrazole (Compound 4) (Scheme II Compound D).

A mixture of (2S)-1-(phenylmethyl)sulfonyl-2-pyrrolidinecarbonitrile (250 mg, 1 mmol), NaN₃ (81 mg, 1.3 mmol) and NH₄Cl (70 mg, 1.3 mmol) in 3 ml DMF was stirred at 130°C for 16 hours. The mixture was concentrated and purified by silica gel to give 120 mg product as a white solid (41.1% yield). ¹H NMR (CDCl₃): δ 01.95 (m, 2H); 2.21 (m, 1H); 2.90 (m, 1H); 3.40 (m, 2H); 4.27 (s, 2H); 5.04 (m, 1H); 7.36-7.41 (m, 5H, aromatic); 8.05 (s, 1H, NH).

Example 6

A lotion comprising the following composition may be

prepared.

	(%)
95% Ethanol	80.0
an N-linked sulfonamide of N-heterocyclic carboxylic acid or carboxylic acid isostere	10.0
α -Tocopherol acetate	0.01
Ethylene oxide (40 mole) adducts of hardened castor oil	0.5
purified water	9.0
perfume and dye	q.s.

Into 95% ethanol are added an N-linked sulfonamides
of N-heterocyclic carboxylic acid or carboxylic acid
isostere, α -tocopherol acetate, ethylene oxide (40
mole) adducts of hardened castor oil, perfume and a
dye. The resulting mixture is stirred and dissolved,
and purified water is added to the mixture to obtain a
transparent liquid lotion.

5 ml of the lotion may be applied once or twice per
day to a site having marked baldness or alopecia.

Example 7

A lotion comprising the following composition shown
may be prepared.

	(%)
95% Ethanol	80.0
an N-linked sulfonamide of N-heterocyclic carboxylic acid or carboxylic acid isostere	0.005
Hinokitol	0.01
Ethylene oxide (40 mole) adducts of hardened castor oil	0.5
Purified water	19.0
Perfume and dye	q.s.

Into 95% ethanol are added an N-linked sulfonamides of N-heterocyclic carboxylic acid or carboxylic acid isostere, hinokitol, ethylene oxide (40 mole) adducts of hardened castor oil, perfume, and a dye. The resulting mixture is stirred, and purified water is added to the mixture to obtain a transparent liquid lotion.

The lotion may be applied by spraying once to 4 times per day to a site having marked baldness or alopecia.

Example 8

An emulsion may be prepared from A phase and B phase having the following compositions.

(A phase)	(%)
Whale wax	0.5
Cetanol	2.0
Petrolatum	5.0
Squalane	10.0
Polyoxyethylene (10 mole) monostearate	2.0
Sorbitan monooleate	1.0
an N-linked sulfonamide of N-heterocyclic carboxylic acid or carboxylic acid isostere	0.01
(B phase)	(%)
Glycerine	10.0
Purified water	69.0
Perfume, dye, and preservative	q.s.

The A phase and the B phase are respectively heated and melted and maintained at 80°C. Both phases are then mixed and cooled under stirring to normal temperature to obtain an emulsion.

The emulsion may be applied by spraying once to 4

times per day to a site having marked baldness or alopecia.

Example 9

A cream may be prepared from A phase and B phase having the following compositions.

(A Phase)	(%)
Fluid paraffin	5.0
Cetostearyl alcohol	5.5
Petrolatum	5.5
Glycerine monostearate	33.0
Polyoxyethylene (20 mole) 2-octyldodecyl ether	3.0
Propylparaben	0.3
(B Phase)	(%)
an N-linked sulfonamide of N-heterocyclic carboxylic acid or carboxylic acid isostere	0.8
Glycerine	7.0
Dipropylene glycol	20.0
Polyethylene glycol 4000	5.0
Sodium Hexametaphosphate	0.005
Purified water	44.895

The A phase is heated and melted, and maintained at 70°C. The B phase is added into the A phase and the mixture is stirred to obtain an emulsion. The emulsion is then cooled to obtain a cream.

The cream may be applied once to 4 times per day to a site having marked baldness or alopecia.

Example 10

A liquid comprising the following composition may be prepared.

	(%)
Polyoxyethylene butyl ether	20.0
Ethanol	50.0
an N-linked sulfonamide of N-heterocyclic carboxylic acid or carboxylic acid isostere	0.001
Propylene glycol	5.0
Polyoxyethylene hardened castor oil derivative (ethylene oxide 80 mole adducts)	0.4
Perfume	q.s.
Purified water	q.s.

Into ethanol are added polyoxypropylene butyl ether, propylene glycol, polyoxyethylene hardened castor oil, an N-linked sulfonamide of N-heterocyclic carboxylic acid or carboxylic acid isostere, and perfume. The resulting mixture is stirred, and purified water is added to the mixture to obtain a liquid.

The liquid may be applied once to 4 times per day to a site having marked baldness or alopecia.

Example 11

A shampoo comprising the following composition may be prepared.

	(%)
Sodium laurylsulfate	5.0
Triethanolamine laurylsulfate	5.0
Betaine lauryldimethylaminoacetate	6.0
Ethylene glycol distearate	2.0
Polyethylene glycol	5.0
an N-linked sulfonamide of N-heterocyclic carboxylic acid or carboxylic acid isostere	5.0

Ethanol	2.0
Perfume	0.3
Purified water	69.7

5 Into 69.7 of purified water are added 5.0 g of sodium laurylsulfate, 5.0 g of triethanolamine laurylsulfate, 6.0 g of betaine lauryldimethyl-aminoacetate. Then a mixture obtained by adding 5.0 g of an N-linked sulfonamide of N-heterocyclic carboxylic acid or
10 carboxylic acid isostere, 5.0 g of polyethylene glycol, and 2.0 g of ethylene glycol distearate to 2.0 g of ethanol, followed by stirring, and 0.3 g of perfume are successively added. The resulting mixture is heated and subsequently cooled to obtain a shampoo.

15 The shampoo may be used on the scalp once or twice per day.

Example 12

20 A patient is suffering from alopecia senilis. An N-linked sulfonamide of N-heterocyclic carboxylic acid or carboxylic acid isostere, or a pharmaceutical composition comprising the same, may be administered to the patient. Increased hair growth is expected to occur following treatment.

Example 13

25 A patient is suffering from male pattern alopecia. An N-linked sulfonamide of N-heterocyclic carboxylic acid or carboxylic acid isostere, or a pharmaceutical composition comprising the same may be administered to the patient. Increased hair growth is expected to occur
30 following treatment.

Example 14

 A patient is suffering from alopecia areata. An N-linked sulfonamide of N-heterocyclic carboxylic acid or

carboxylic acid isostere, or a pharmaceutical composition comprising the same, may be administered to the patient. Increased hair growth is expected to occur following treatment.

5

Example 15

A patient is suffering from hair loss caused by skin lesions. An N-linked sulfonamide of N-heterocyclic carboxylic acid or carboxylic acid isostere, or a pharmaceutical composition comprising the same, may be administered to the patient. Increased hair growth is expected to occur following treatment.

10

Example 16

A patient is suffering from hair loss caused by tumors. An N-linked sulfonamide of N-heterocyclic carboxylic acid or carboxylic acid isostere, or a pharmaceutical composition comprising the same, may be administered to the patient. Increased hair growth is expected to occur following treatment.

15

Example 17

A patient is suffering from hair loss caused by a systematic disorder, such as a nutritional disorder or an internal secretion disorder. An N-linked sulfonamide of N-heterocyclic carboxylic acid or carboxylic acid isostere, or a pharmaceutical composition comprising the same, may be administered to the patient. Increased hair growth is expected to occur following treatment.

20

25

Example 18

A patient is suffering from hair loss caused by chemotherapy. An N-linked sulfonamide of N-heterocyclic carboxylic acid or carboxylic acid isostere, or a pharmaceutical composition comprising the same, may be administered to the patient. Increased hair growth is expected to occur following treatment.

30

Example 19

5 A patient is suffering from hair loss caused by radiation. An N-linked sulfonamide of N-heterocyclic carboxylic acid or carboxylic acid isostere, or a pharmaceutical composition comprising the same may, be administered to the patient. Increased hair growth is expected to occur following treatment.

Example 20

10 A patient is suffering from a neurodegenerative disease. A carboxylic acid or carboxylic acid isostere of an N-heterocyclic ring or a pharmaceutical composition comprising the same is administered. It would be expected that the patient would improve their condition or recover.

Example 21

15 A patient is suffering from a neurological disorder. A carboxylic acid or carboxylic acid isostere of an N-heterocyclic ring or pharmaceutical compositions comprising same is administered. It would be expected that the patient would improve their condition or recover.

Example 22

25 A patient is suffering from stroke. A carboxylic acid or carboxylic acid isostere of an N-heterocyclic ring or pharmaceutical compositions comprising same is administered. It would be expected that the patient would improve their condition or recover.

Example 23

30 A patient is suffering from Parkinson's Disease. A carboxylic acid or carboxylic acid isostere of an N-heterocyclic ring or pharmaceutical compositions comprising same is administered. It would be expected that the patient would improve their condition or recover.

Example 24

5 A patient is suffering from Alzheimer's Disease. A carboxylic acid or carboxylic acid isostere of an N-heterocyclic ring or pharmaceutical compositions comprising same is administered. It would be expected that the patient would improve their condition or recover.

Example 25

10 A patient is suffering from a peripheral neuropathy. A carboxylic acid or carboxylic acid isostere of an N-heterocyclic ring or pharmaceutical compositions comprising same is administered. It would be expected that the patient would improve their condition or recover.

Example 26

15 A patient is suffering from amyotrophic lateral sclerosis. A carboxylic acid or carboxylic acid isostere of an N-heterocyclic ring or pharmaceutical compositions comprising same is administered. It would be expected that the patient would improve their condition or recover.

Example 27

25 A patient is suffering from a spinal injury. A carboxylic acid or carboxylic acid isostere of an N-heterocyclic ring or pharmaceutical compositions comprising same is administered. It would be expected that the patient would improve their condition or recover.

Example 28


30 A patient is at risk of suffering from a neurodegenerative disease or neurological disorder. A carboxylic acid or carboxylic acid isostere of an N-heterocyclic ring or a pharmaceutical composition comprising the same is prophelactically administered.

It would be expected that the patient would be prevented from some or all of the effects of the disease or disorder, or would significantly improve their condition or recover over patients who were not pre-treated.

5 The invention being thus described, it will be obvious
that the same may be varied in many ways. Such
variations are not to be regarded as a departure from
the spirit and scope of the invention and all such
modification are intended to be included within the
10 scope of the following claims.

What is claimed is:

1. A compound having the formula (I):



where

n is 1-3;

R_1 is selected from the group consisting of hydrogen, C_1 - C_9 straight or branched chain alkyl, C_2 - C_9 straight or branched chain alkenyl, aryl, heteroaryl, carbocycle, or heterocycle;

D is a bond, or a C_1 - C_{10} straight or branched chain alkyl, C_2 - C_{10} alkenyl or C_2 - C_{10} alkynyl;

R_2 is a carboxylic acid or a carboxylic acid isostere;

wherein said alkyl, alkenyl, alkynyl, aryl, heteroaryl, carbocycle, heterocycle, or carboxylic acid isostere is optionally substituted with one or more substituents selected from R^3 , where

R^3 is hydrogen, hydroxy, halo, haloalkyl, thiocarbonyl, alkoxy, alkenoxy, alkylaryloxy, aryloxy, arylalkyloxy, cyano, nitro, imino, alkylamino, aminoalkyl, sulfhydryl, thioalkyl, alkylthio, sulfonyl, C_1 - C_6 straight or branched chain alkyl, C_2 - C_6 straight or branched chain alkenyl or alkynyl, aryl, heteroaryl, carbocycle, heterocycle, or CO_2R^4 where R^4 is hydrogen or C_1 - C_9 straight or branched chain alkyl or alkenyl;

or a pharmaceutically acceptable salt, ester or solvate thereof;

provided that:

when D is a bond, and R₂ is COOH,
then R₁ cannot be substituted naphthyl;
further provided that:

when D is a bond, and n is 1, and R₂ is COOH or CONHR₃,
then R₁ is not hydroxyl, methyl, ethyl, substituted or
unsubstituted thioethyl, benzothiazole, substituted
benzopyran, substituted benzopyrrole, substituted
benzoxazole, substituted 5-membered heterocycle
containing two N and one S heteroatoms, or substituted
or unsubstituted phenyl, phenylethyl, naphthyl, pyridyl,
thienyl, quinoline, tricyclic ring, aminoethyl, or
benzyl;

further provided that:

when D is a bond, and n is 2, and R₂ is COOH or
phenylbutyl ester,
then R₁ is not substituted phenyl, or a substituted
bicyclic ring containing two oxygen heteroatoms.

further provided that:

when D is a bond, and n is 1-2, and R₂ is a substituted
or unsubstituted carbocyclic or heterocyclic ring
structure,
then R₁ is not substituted or unsubstituted carbocycle
or heterocycle, or hydroxy;

further provided that:

when D is a bond, and n is 1-2, and R₂ is hydroxy,
alkoxy, -SO₂(phenyl), N(R₃)₂, substituted thio or
alkylthio, -NCO, -PO₃(Me)₂, or -NCOOC(ethyl)phenyl,
then R₁ is not naphthalene, ethylene, substituted
tricyclic ring, or substituted or unsubstituted phenyl;

further provided that:

when D is C₁-C₃ alkyl or hexenyl, and R₂ is hydroxyl,
then R₁ is not substituted or unsubstituted phenyl, or
benzoimidazole;

further provided that:

when D is methyl, and n is 1, and R₂ is cyano or COOH,

then R₁ is not substituted phenyl;

further provided that:

when D is methyl, and n is 1, and R₂ is methoxy or N(R₃)₂,

5 then R₁ is not methyl, ethyl, phenylethyl, chloro substituted alkyl, substituted oxirane, substituted aziridine wherein one of the carbons is replaced with an oxygen, substituted or unsubstituted propenyl, substituted phenyl, benzyl, or trifluoro substituted C₂-
10 C₃ alkyl or alkenyl;

further provided that:

when D is ethyl, and n is 2, and R₂ is hydroxyl or N(R₃)₂,

then R₁ is not naphthyl;

15 further provided that:

when D is propyl, and n is 1, and R₂ is methoxy,

then R₁ is not ethylene, cyano substituted ethyl, or triethoxy substituted propyl;


further provided that:

20 when D is not a bond and at least one of D and R₂ contains at least one S or O,

then R₁ is not methyl or substituted phenyl.

25 2. The compound of claim 1, wherein R₂ is a carbocycle or heterocycle containing any combination of CH₂, O, S, or N in any chemically stable oxidation state, wherein any of the atoms of said ring structure are optionally substituted in one or more positions with R³.

30 3. The compound of claim 1, wherein R₂ is selected from the following group:



where the atoms of said ring structure may be optionally substituted at one or more positions with R^3 .

4. The compound of claim 1, wherein the carboxylic acid or carboxylic acid isostere of R_2 is selected from the group consisting of:


$-\text{COOH}$, $-\text{SO}_3\text{H}$, $-\text{SO}_2\text{HNR}^3$, $-\text{PO}_2(\text{R}^3)_2$, $-\text{CN}$, $-\text{PO}_3(\text{R}^3)_2$, $-\text{OR}^3$, $-\text{SR}^3$, $-\text{NHCOR}^3$, $-\text{N}(\text{R}^3)_2$, $-\text{CON}(\text{R}^3)_2$, $-\text{CONH}(\text{O})\text{R}^3$, $-\text{CONHNHSO}_2\text{R}^3$, $-\text{COHNSO}_2\text{R}^3$, and $-\text{CONR}^3\text{CN}$.

5. The compounds, (2S)-1-(phenylmethyl)sulfonyl-2-hydroxymethyl pyrrolidine; (2S)-1-(phenylmethyl)sulfonyl-2-pyrrolidinetetrazole.

6. A pharmaceutical composition, comprising:

- a) an effective amount of an N-linked sulfonamide of an N-heterocyclic carboxylic acid or carboxylic acid isostere; and
- b) a pharmaceutically acceptable carrier.

7. The pharmaceutical composition of claim 6, wherein the N-linked sulfonamide of N-heterocyclic carboxylic acid or carboxylic acid isostere comprises a compound of formula (I):



where

n is 1-3;

R_1 is selected from the group consisting of hydrogen, C_1-C_9 straight or branched chain alkyl, C_2-C_9 straight or branched chain alkenyl, aryl, heteroaryl, carbocycle, or heterocycle;

5 D is a bond, or a C_1-C_{10} straight or branched chain alkyl, C_2-C_{10} alkenyl or C_2-C_{10} alkynyl;


R_2 is a carboxylic acid or a carboxylic acid isostere; wherein said alkyl, alkenyl, alkynyl, aryl, heteroaryl, carbocycle, heterocycle, or carboxylic acid isostere is optionally substituted with one or more substituents selected from R^3 , where

10 R^3 is hydrogen, hydroxy, halo, haloalkyl, thiocarbonyl, alkoxy, alkenoxy, alkylaryloxy, aryloxy, arylalkyloxy, cyano, nitro, imino, alkylamino, aminoalkyl, sulfhydryl, thioalkyl, alkylthio, sulfonyl, C_1-C_6 straight or branched chain alkyl, C_2-C_6 straight or branched chain alkenyl or alkynyl, aryl, heteroaryl, carbocycle, heterocycle, or CO_2R^4 where R^4 is hydrogen or C_1-C_9 straight or branched chain alkyl or alkenyl;

15 or a pharmaceutically acceptable salt, ester or solvate thereof.

8. The pharmaceutical composition of claim 7, wherein R_2 is a carbocycle or heterocycle containing any combination of CH_2 , O, S, or N in any chemically stable oxidation state, wherein any of the atoms of said ring structure are optionally substituted in one or more positions with R^3 .

30 9. The pharmaceutical composition of claim 7, wherein R_2 is selected from the following group:



where the atoms of said ring structure may be optionally substituted at one or more positions with R³.

10. The pharmaceutical composition of claim 7, wherein
5 R₂ is selected from the group consisting of:
-COOH, -SO₃H, -SO₂HNR³, -PO₂(R³)₂, -CN, -PO₃(R³)₂, -OR³, -
SR³, -NHCOR³, -N(R³)₂, -CON(R³)₂, -CONH(O)R³,
-CONHNHSO₂R³, -COHNSO₂R³, and -CONR³CN.

10 11. The pharmaceutical composition of claim 7, wherein
the N-linked sulfonamide of an N-heterocyclic
carboxylic acid compound is selected from the group
consisting of compounds 1-97.

15 12. The pharmaceutical composition of claim 6, further
comprising a neurotrophic factor different from formula
(I).

20 13. The pharmaceutical composition of claim 12, wherein
said neurotrophic factor different from formula (I) is
selected from neurotrophic growth factor, brain derived
growth factor, glial derived growth factor, ciliary
neurotrophic factor, insulin growth factor and active
truncated derivatives thereof, acidic fibroblast growth
25 factor, basic fibroblast growth factor, platelet-derived
growth factors, neurotrophin-3 and neurotrophin 4/5.

14. A method of treating a neurological disorder in an
animal, comprising:

30 administering to the animal an effective amount of an
N-linked sulfonamide of an N-heterocyclic carboxylic
acid or carboxylic acid isostere to stimulate growth of
damaged peripheral nerves or to promote neuronal
regeneration.

15. The method of claim 14, wherein the neurological disorder is selected from the group consisting of peripheral neuropathies cause by physical injury or disease state, physical damage to the brain, physical damage to the spinal cord, stroke associated with brain damage, and neurological disorders relating to neurodegeneration.

16. The method of claim 14, wherein the neurological disorder is selected from the group consisting of Alzheimer's Disease, Parkinson's Disease, and amyotrophic lateral sclerosis.


17. The method of claim 14, wherein the neurological disorder is Alzheimer's disease.

18. The method of claim 14, wherein the neurological disorder is Parkinson's disease.

19. The method of claim 14, wherein the neurological disorder is amyotrophic lateral sclerosis.

20. The method of claim 14, wherein the N-linked sulfonamide of an N-heterocyclic carboxylic acid or carboxylic acid isostere is non-immunosuppressive.

21. The method of claim 14, wherein the N-linked sulfonamide of an N-heterocyclic carboxylic acid or carboxylic acid isostere comprises a compound of formula (I):



where

n is 1-3;

5 R₁ is selected from the group consisting of hydrogen, C₁-C₉ straight or branched chain alkyl, C₂-C₉ straight or branched chain alkenyl, aryl, heteroaryl, carbocycle, or heterocycle;

D is a bond, or a C₁-C₁₀ straight or branched chain alkyl, C₂-C₁₀ alkenyl or C₂-C₁₀ alkynyl;

10 R₂ is a carboxylic acid or a carboxylic acid isostere; wherein said alkyl, alkenyl, alkynyl, aryl, heteroaryl, carbocycle, heterocycle, or carboxylic acid isostere is optionally substituted with one or more substituents selected from R³, where

15 R³ is hydrogen, hydroxy, halo, haloalkyl, thiocarbonyl, alkoxy, alkenoxy, alkylaryloxy, aryloxy, arylalkyloxy, cyano, nitro, imino, alkylamino, aminoalkyl, sulfhydryl, thioalkyl, alkylthio, sulfonyl, C₁-C₆ straight or branched chain alkyl, C₂-C₆ straight or branched chain alkenyl or alkynyl, aryl, heteroaryl, carbocycle, heterocycle, or CO₂R⁴ where R⁴ is hydrogen or C₁-C₉ straight or branched chain alkyl or alkenyl;

20 or a pharmaceutically acceptable salt, ester or solvate thereof.


25

22. The method of claim 21, wherein R₂ is a carbocycle or heterocycle containing any combination of CH₂, O, S, or N in any chemically stable oxidation state, wherein

any of the atoms of said ring structure are optionally substituted in one or more positions with R³.

5

23. The method of claim 21, wherein R₂ is selected from the following group:



where the atoms of said ring structure may be optionally substituted at one or more positions with R³.

5 24. The method of claim 21, wherein R₂ is selected from the group consisting of:

-COOH, -SO₃H, -SO₂HNR³, -PO₂(R³)₂, -CN, -PO₃(R³)₂, -OR³, -SR³, -NHCOR³, -N(R³)₂, -CON(R³)₂, -CONH(O)R³, -CONHNHSO₂R³, -COHNSO₂R³, and -CONR³CN.

10 25. The method of claim 14, wherein the N-linked sulfonamide of an N-heterocyclic carboxylic acid compound is selected from the group consisting of compounds 1-97.

15 26. The method of claim 14, further comprising administering a neurotrophic factor different from formula (I).

20 27. The method of claim 26, wherein said neurotrophic factor different from formula (I) is selected from the group consisting of neurotrophic growth factor, brain derived growth factor, glial derived growth factor, ciliary neurotrophic factor, insulin growth factor and active truncated derivatives thereof, acidic fibroblast growth factor, basic fibroblast growth factor, platelet-derived growth factors, neurotrophin-3, and neurotrophin 4/5.


30 28. A method of stimulating growth of damaged peripheral nerves, comprising:

administering to damaged peripheral nerves a therapeutically effective amount of an N-linked sulfonamide of an N-heterocyclic carboxylic acid or
35 carboxylic acid isostere to stimulate or promote growth

of the damaged peripheral nerves.

29. The method of claim 28, wherein the N-linked sulfonamide of an N-heterocyclic carboxylic acid or carboxylic acid isostere is non-immunosuppressive.

30. The method of claim 28, wherein the N-linked sulfonamide of an N-heterocyclic carboxylic acid or carboxylic acid isostere comprises a compound of formula (I):



where

n is 1-3;

R₁ is selected from the group consisting of hydrogen, C₁-C₉ straight or branched chain alkyl, C₂-C₉ straight or branched chain alkenyl, aryl, heteroaryl, carbocycle, or heterocycle;

D is a bond, or a C₁-C₁₀ straight or branched chain alkyl, C₂-C₁₀ alkenyl or C₂-C₁₀ alkynyl;

R₂ is a carboxylic acid or a carboxylic acid isostere; wherein said alkyl, alkenyl, alkynyl, aryl, heteroaryl, carbocycle, heterocycle, or carboxylic acid isostere is optionally substituted with one or more substituents selected from R³, where


R³ is hydrogen, hydroxy, halo, haloalkyl, thiocarbonyl, alkoxy, alkenoxy, alkylaryloxy, aryloxy, arylalkyloxy, cyano, nitro, imino, alkylamino, aminoalkyl, sulfhydryl, thioalkyl, alkylthio, sulfonyl, C₁-C₆ straight or

branched chain alkyl, C₂-C₆ straight or branched chain alkenyl or alkynyl, aryl, heteroaryl, carbocycle, heterocycle, or CO₂R⁴ where R⁴ is hydrogen or C₁-C₉ straight or branched chain alkyl or alkenyl;

5 or a pharmaceutically acceptable salt, ester or solvate thereof.

31. The method of claim 30, wherein R₂ is a carbocycle or heterocycle containing any combination of CH₂, O, S, or N in any chemically stable oxidation state, wherein
10 any of the atoms of said ring structure are optionally substituted in one or more positions with R³.

32. The method of claim 30, wherein R₂ is selected from
15 the following group:



where the atoms of said ring structure may be optionally substituted at one or more positions with R³.

5 33. The method of claim 30, wherein R₂ is selected from the group consisting of:

-COOH, -SO₃H, -SO₂HNR³, -PO₂(R³)₂, -CN, -PO₃(R³)₂, -OR³, -SR³, -NHCOR³, -N(R³)₂, -CON(R³)₂, -CONH(O)R³, -CONHNHSO₂R³, -COHNSO₂R³, and -CONR³CN.

10 34. The method of claim 28, wherein the N-linked sulfonamide of an N-heterocyclic carboxylic acid compound is selected from the group consisting of compounds 1-97.

15 35. The method of claim 28, further comprising administering a neurotrophic factor different from formula (I).

20 36. The method of claim 35, wherein said neurotrophic factor different from formula (I) is selected from the group consisting of neurotrophic growth factor, brain derived growth factor, glial derived growth factor, ciliary neurotrophic factor, insulin growth factor and active truncated derivatives thereof, acidic fibroblast growth factor, basic fibroblast growth factor, platelet-derived growth factors, neurotrophin-3, and neurotrophin 4/5.


30 37. A method for promoting neuronal regeneration and growth in animals, comprising:

administering to an animal a therapeutically effective amount of a neurotrophic N-linked sulfonamide of an N-heterocyclic carboxylic acid or carboxylic acid isostere to promote neuronal regeneration.

35

38. The method of claim 37, wherein the N-linked sulfonamide of an N-heterocyclic carboxylic acid or carboxylic acid isostere is non-immunosuppressive.

5 39. The method of claim 37, wherein the N-linked sulfonamide of an N-heterocyclic carboxylic acid or carboxylic acid isostere comprises a compound of formula (I):



10 where

n is 1-3;

R₁ is selected from the group consisting of hydrogen, C₁-C₉ straight or branched chain alkyl, C₂-C₉ straight or branched chain alkenyl, aryl, heteroaryl, carbocycle, or heterocycle;

D is a bond, or a C₁-C₁₀ straight or branched chain alkyl, C₂-C₁₀ alkenyl or C₂-C₁₀ alkynyl;

R₂ is a carboxylic acid or a carboxylic acid isostere;

15 wherein said alkyl, alkenyl, alkynyl, aryl, heteroaryl, carbocycle, heterocycle, or carboxylic acid isostere is optionally substituted with one or more substituents selected from R³, where

R³ is hydrogen, hydroxy, halo, haloalkyl, thiocarbonyl, alkoxy, alkenoxy, alkylaryloxy, aryloxy, arylalkyloxy, cyano, nitro, imino, alkylamino, aminoalkyl, sulfhydryl, thioalkyl, alkylthio, sulfonyl, C₁-C₆ straight or branched chain alkyl, C₂-C₆ straight or branched chain alkenyl or alkynyl, aryl, heteroaryl, carbocycle,

25




















heterocycle, or CO_2R^4 where R^4 is hydrogen or $\text{C}_1\text{-C}_6$,
straight or branched chain alkyl or alkenyl;
or a pharmaceutically acceptable salt, ester or solvate
thereof.

5

40. The method of claim 39, wherein R_2 is a carbocycle
or heterocycle containing any combination of CH_2 , O, S,
or N in any chemically stable oxidation state, wherein
any of the atoms of said ring structure are optionally
substituted in one or more positions with R^3 .

10

41. The method of claim 39, wherein R_2 is selected from
the following group:



where the atoms of said ring structure may be optionally substituted at one or more positions with R³.

5 42. The method of claim 39, wherein R₂ is selected from the group consisting of:

-COOH, -SO₃H, -SO₂HNR³, -PO₂(R³)₂, -CN, -PO₃(R³)₂, -OR³, -SR³, -NHCOR³, -N(R³)₂, -CON(R³)₂, -CONH(O)R³, -CONHNHSO₂R³, -COHNSO₂R³, and -CONR³CN.

10 43. The method of claim 37, wherein the N-linked sulfonamide of an N-heterocyclic carboxylic acid compound is selected from the group consisting of compounds 1-97.

15 44. The method of claim 37, further comprising administering a neurotrophic factor different from formula (I).

20 45. The method of claim 44, wherein said neurotrophic factor different from formula (I) is selected from the group consisting of neurotrophic growth factor, brain derived growth factor, glial derived growth factor, ciliary neurotrophic factor, insulin growth factor and active truncated derivatives thereof, acidic fibroblast growth factor, basic fibroblast growth factor, platelet-derived growth factors, neurotrophin-3, and neurotrophin 4/5.

30 46. A method for preventing neurodegeneration in an animal, comprising:

administering to an animal a therapeutically effective amount of a neurotrophic N-linked sulfonamide of an N-heterocyclic carboxylic acid or carboxylic acid isostere to prevent neurodegeneration.

35


47. The method of claim 46, wherein the neurodegeneration is Alzheimer's disease.

48. The method of claim 46, wherein the neurodegeneration is Parkinson's disease.

49. The method of claim 46, wherein the neurodegeneration is amyotrophic lateral sclerosis.

50. The method of claim 46, wherein the N-linked sulfonamide of an N-heterocyclic carboxylic acid or carboxylic acid isostere is non-immunosuppressive.

51. The method of claim 46, wherein the N-linked sulfonamide of an N-heterocyclic carboxylic acid or carboxylic acid isostere comprises a compound of formula (I):



where

n is 1-3;

R₁ is selected from the group consisting of hydrogen, C₁-C₉ straight or branched chain alkyl, C₂-C₉ straight or branched chain alkenyl, aryl, heteroaryl, carbocycle, or heterocycle;

D is a bond, or a C₁-C₁₀ straight or branched chain alkyl, C₂-C₁₀ alkenyl or C₂-C₁₀ alkynyl;

R₂ is a carboxylic acid or a carboxylic acid isostere;


wherein said alkyl, alkenyl, alkynyl, aryl, heteroaryl,

carbocycle, heterocycle, or carboxylic acid isostere is optionally substituted with one or more substituents selected from R^3 , where

5 R^3 is hydrogen, hydroxy, halo, haloalkyl, thiocarbonyl, alkoxy, alkenoxy, alkylaryloxy, aryloxy, arylalkyloxy, cyano, nitro, imino, alkylamino, aminoalkyl, sulfhydryl, thioalkyl, alkylthio, sulfonyl, C_1 - C_6 straight or
10 branched chain alkyl, C_2 - C_6 straight or branched chain alkenyl or alkynyl, aryl, heteroaryl, carbocycle, heterocycle, or CO_2R^4 where R^4 is hydrogen or C_1 - C_6 straight or branched chain alkyl or alkenyl;
or a pharmaceutically acceptable salt, ester or solvate thereof.

15 52. The method of claim 51, wherein R_2 is a carbocycle or heterocycle containing any combination of CH_2 , O, S, or N in any chemically stable oxidation state, wherein any of the atoms of said ring structure are optionally substituted in one or more positions with R^3 .

20 53. The method of claim 51, wherein R_2 is selected from the following group:



where the atoms of said ring structure may be optionally substituted at one or more positions with R³.

5 54. The method of claim 51, wherein R₂ is selected from the group consisting of:

-COOH, -SO₃H, -SO₂HNR³, -PO₂(R³)₂, -CN, -PO₃(R³)₂, -OR³, -SR³, -NHCOR³, -N(R³)₂, -CON(R³)₂, -CONH(O)R³, -CONHNHSO₂R³, -COHNSO₂R³, and -CONR³CN.

10 55. The method of claim 46, wherein the N-linked sulfonamide of an N-heterocyclic carboxylic acid compound is selected from the group consisting of compounds 1-97.

15 56. The method of claim 46, further comprising administering a neurotrophic factor different from formula (I).


20 57. The method of claim 56, wherein said neurotrophic factor different from formula (I) is selected from the group consisting of neurotrophic growth factor, brain derived growth factor, glial derived growth factor, ciliary neurotrophic factor, insulin growth factor and active truncated derivatives thereof, acidic fibroblast growth factor, basic fibroblast growth factor, platelet-derived growth factors, neurotrophin-3, and neurotrophin 4/5.

30 58. A method for treating alopecia or promoting hair growth in an animal, which comprises administering to said animal an effective amount of an N-linked sulfonamide of an N-heterocyclic carboxylic acid or carboxylic acid isostere.

35

59. The method of claim 58, wherein the N-linked sulfonamide of an N-heterocyclic carboxylic acid or carboxylic acid isostere is non-immunosuppressive.

5 60. The method of claim 58, wherein the N-linked sulfonamide of an N-heterocyclic carboxylic acid or carboxylic acid isostere is a compound of formula (I):



where

10 n is 1-3;

R₁ is selected from the group consisting of hydrogen, C₁-C₉ straight or branched chain alkyl, C₂-C₉ straight or branched chain alkenyl, aryl, heteroaryl, carbocycle, or heterocycle;

15 D is a bond, or a C₁-C₁₀ straight or branched chain alkyl, C₂-C₁₀ alkenyl or C₂-C₁₀ alkynyl;

R₂ is a carboxylic acid or a carboxylic acid isostere;

20 wherein said alkyl, alkenyl, alkynyl, aryl, heteroaryl, carbocycle, heterocycle, or carboxylic acid isostere is optionally substituted with one or more substituents selected from R³, where

R³ is hydrogen, hydroxy, halo, haloalkyl, thiocarbonyl, alkoxy, alkenoxy, alkylaryloxy, aryloxy, arylalkyloxy, cyano, nitro, imino, alkylamino, aminoalkyl, sulfhydryl, thioalkyl, alkylthio, sulfonyl, C₁-C₆ straight or branched chain alkyl, C₂-C₆ straight or branched chain alkenyl or alkynyl, aryl, heteroaryl, carbocycle, heterocycle, or CO₂R⁴ where R⁴ is hydrogen or C₁-C₉,


25

straight or branched chain alkyl or alkenyl;
or a pharmaceutically acceptable salt, ester or solvate
thereof.

5 61. The method of claim 60, wherein R_2 is a carbocycle
or heterocycle containing any combination of CH_2 , O, S,
or N in any chemically stable oxidation state, wherein
any of the atoms of said ring structure are optionally
substituted in one or more positions with R^3 .

10

62. The method of claim 60, wherein R_2 is selected from
the following group:



where the atoms of said ring structure may be optionally substituted at one or more positions with R^3 .

63. The method of claim 60, wherein R_2 is selected from the group consisting of
 $-\text{COOH}$, $-\text{SO}_3\text{H}$, $-\text{SO}_2\text{HNR}^3$, $-\text{PO}_2(\text{R}^3)_2$, $-\text{CN}$, $-\text{PO}_3(\text{R}^3)_2$, $-\text{OR}^3$, $-\text{SR}^3$, $-\text{NHCOR}^3$, $-\text{N}(\text{R}^3)_2$, $-\text{CON}(\text{R}^3)_2$, $-\text{CONH}(\text{O})\text{R}^3$, $-\text{CONHNHSO}_2\text{R}^3$, $-\text{COHNSO}_2\text{R}^3$, and $-\text{CONR}^3\text{CN}$.


64. The method of claim 58, wherein the carboxylic acid or carboxylic acid isostere is selected from the group consisting of compounds 1-97.

65. A pharmaceutical composition comprising:

- (i) an effective amount of a N-linked sulfonamide of an N-heterocyclic carboxylic acid or carboxylic acid isostere for treating alopecia or promoting hair growth in an animal; and
- (ii) a pharmaceutically acceptable carrier.

66. The pharmaceutical composition of claim 65, wherein the N-linked sulfonamide of an N-heterocyclic carboxylic acid or carboxylic acid isostere is non-immunosuppressive.

67. The composition of claim 65, wherein the carboxylic acid or carboxylic acid isostere is a compound of formula (I):



where

n is 1-3;

R₁ is selected from the group consisting of hydrogen, C₁-C₉ straight or branched chain alkyl, C₂-C₉ straight or branched chain alkenyl, aryl, heteroaryl, carbocycle, or heterocycle;

D is a bond, or a C₁-C₁₀ straight or branched chain alkyl, C₂-C₁₀ alkenyl or C₂-C₁₀ alkynyl;

R₂ is a carboxylic acid or a carboxylic acid isostere;


wherein said alkyl, alkenyl, alkynyl, aryl, heteroaryl, carbocycle, heterocycle, or carboxylic acid isostere is optionally substituted with one or more substituents selected from R³, where

R³ is hydrogen, hydroxy, halo, haloalkyl, thiocarbonyl, alkoxy, alkenoxy, alkylaryloxy, aryloxy, arylalkyloxy, cyano, nitro, imino, alkylamino, aminoalkyl, sulfhydryl, thioalkyl, alkylthio, sulfonyl, C₁-C₆ straight or branched chain alkyl, C₂-C₆ straight or branched chain alkenyl or alkynyl, aryl, heteroaryl, carbocycle, heterocycle, or CO₂R⁴ where R⁴ is hydrogen or C₁-C₉ straight or branched chain alkyl or alkenyl;

or a pharmaceutically acceptable salt, ester or solvate thereof.

68. The composition of claim 67, wherein R₂ is a carbocycle or heterocycle containing any combination of CH₂, O, S, or N in any chemically stable oxidation state, wherein any of the atoms of said ring structure are optionally substituted in one or more positions with R³.

69. The composition of claim 67, wherein R₂ is selected from the following group:



where the atoms of said ring structure may be optionally substituted at one or more positions with R^3 .

5 70. The composition of claim 67, wherein R_2 is selected from the group consisting of:

-COOH, -SO₃H, -SO₂HNR³, -PO₂(R³)₂, -CN, -PO₃(R³)₂, -OR³, -SR³, -NHCOR³, -N(R³)₂, -CON(R³)₂, -CONH(O)R³, -CONHNHSO₂R³, -COHNSO₂R³, and -CONR³CN.

10 71. The composition of claim 65, wherein the carboxylic acid or carboxylic acid isostere is selected from the group consisting of compounds 1-97.

FIG. 1




FIG.2





FIG. 3

Promotion of Hair Growth by Neuroimmunophilin Ligands



INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/25572

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D207/16 A61K31/40 A61K31/41 C07D207/08 C07D403/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 614 547 A (HAMILTON GREGORY S ET AL) 25 March 1997 see abstract; claims see column 13 - column 21; examples ---	1-71
X	EP 0 610 744 A (MILES INC) 17 August 1994 see abstract; claim 1; examples 1-9 see page 3, line 52 - line 57 ---	1-71
X	WO 92 21313 A (VERTEX PHARMA) 10 December 1992 see abstract; claims see page 12 - page 13; tables see page 22 - page 25; example 2 --- -/--	1-11

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"8" document member of the same patent family

Date of the actual completion of the international search

22 April 1999

Date of mailing of the international search report

04/05/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Paisdor, B

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/25572

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 96 40140 A (GUILFORD PHARM INC ;UNIV JOHNS HOPKINS MED (US)) 19 December 1996 see abstract; claims see page 57; table 2 ---	1-71
X	WO 98 08827 A. (PROCTER & GAMBLE) 5 March 1998 see page 41 - page 45; examples 13E-G, 14C-E see page 47; examples 91-101 see page 48 - page 54; examples see page 17, line 20 - line 24 ---	1-71
E	WO 98 55090 A (GUILFORD PHARM INC) 10 December 1998 see page 94 - page 96 see page 114 see page 165 - page 175; examples ---	1-71
A	KING F D: "BIOISOSTERES, CONFORMATIONAL RESTRICTION, AND PRO-DRUGS - CASE HISTORY: AN EXAMPLE OF A CONFORMATIONAL RESTRICTION APPROACH" 1994 , MEDICINAL CHEMISTRY: PRINCIPLES AND PRACTICE, PAGE(S) 206 - 209 , KING F D XP002033086 see page 208 -----	1

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 98/25572

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 14-64
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 14-64
are directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☒ Claims Nos.: not applicable
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Claims Nos.: not applicable

In view of the extremely broad Markush claims, the search was executed with due regard to the PCT Search Guidelines (PCT/GL/2), C-III, paragraph 2.1, 2.3 read in conjunction with 3.7 and Rule 33.3 PCT, i.e. particular emphasis was put on the inventive concept, as illustrated by claim 5 and the specific embodiments mentioned in table I of the present application, and by those compounds which were actually prepared and for which physical data was given. The international search was, in so far as possible and reasonable, complete in that it covered the entire subject-matter to which the claims are directed.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 98/25572

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5614547	A	25-03-1997	AU 703118 B	18-03-1999
			AU 6106296 A	30-12-1996
			BG 102071 A	30-09-1998
			BR 9608444 A	05-01-1999
			CA 2206799 A	19-12-1996
			CH 688775 A	13-03-1998
			CZ 9702330 A	17-12-1997
			DE 19680256 T	19-06-1997
			DK 125796 A	19-12-1996
			EP 0769006 A	23-04-1997
			FI 964328 A	30-12-1996
			GB 2305176 A	02-04-1997
			GB 2324527 A	28-10-1998
			GB 2325230 A	18-11-1998
			JP 8333334 A	17-12-1996
			LT 98001 A,B	25-11-1998
			LU 88833 A	15-01-1997
			LV 11991 A	20-03-1998
			LV 11991 B	20-07-1998
			NO 974213 A	04-12-1997
			PL 323300 A	16-03-1998
			SE 9604098 A	08-12-1996
			SI 9620091 A	31-12-1998
			SK 158597 A	09-09-1998
			WO 9640633 A	19-12-1996
			US 5859031 A	12-01-1999
			US 5795908 A	18-08-1998
EP 0610744	A	17-08-1994	AT 172985 T	15-11-1998
			CA 2115222 A	10-08-1994
			DE 69414279 D	10-12-1998
			DE 69414279 T	08-04-1999
			ES 2124327 T	01-02-1999
			JP 6234735 A	23-08-1994
			US 5633277 A	27-05-1997
WO 9221313	A	10-12-1992	AU 2185092 A	08-01-1993
			CA 2102178 A	25-11-1992
			EP 0587756 A	23-03-1994
			JP 6508141 T	14-09-1994
			MX 9202466 A	30-06-1994
WO 9640140	A	19-12-1996	US 5798355 A	25-08-1998
			US 5696135 A	09-12-1997
			AU 6162296 A	30-12-1996
			BG 102072 A	31-08-1998
			CA 2206824 A	19-12-1996
			CZ 9702329 A	15-07-1998
			DE 19680255 T	05-06-1997
			DK 125696 A	20-12-1996
			EP 0777478 A	11-06-1997
			FI 964137 A	15-01-1997
			GB 2305605 A	16-04-1997
			JP 8333256 A	17-12-1996
			LT 98002 A	25-02-1999
			LU 88834 A	15-01-1997
			LV 11986 A	20-03-1998
			LV 11986 B	20-09-1998

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 98/25572

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9640140 A		NO 974290 A	04-12-1997
		PL 323381 A	30-03-1998
		SE 9604097 A	08-12-1997
		US 5843960 A	01-12-1998
		US 5846981 A	08-12-1998
WO 9808827 A	05-03-1998	AU 4073997 A	19-03-1998
		AU 4152997 A	19-03-1998
		WO 9808825 A	05-03-1998
WO 9855090 A	10-12-1998	AU 7716498 A	21-12-1998
		AU 7716998 A	21-12-1998
		WO 9855091 A	10-12-1998